

Mechanisms of Thermoregulation

207

The process of controlling body temperature is called thermoregulation. For many years, animals were classified as either homeotherms or homoiotherms (= constant body temperature) or poikilotherms (= variable body temperature). Unfortunately, these terms are not particularly accurate for many animals; for example, some mammals (typical homeotherms),

may have unstable body temperatures. A more recent, thermal classification of animals is based on the source of the body heat: whether it is largely from the environment (ectothermic) or from metabolic activity (endothermic). This classification can be more accurately applied to most animals but, in reality, many animals still fall somewhere between the two extremes.

How Body Temperature Varies




Aquatic invertebrates like jellyfish are true poikilotherms: their temperature is the same as the environment.



Tuna and some of the larger sharks can maintain body temperatures up to 14°C above the water temperature.



Hibernating rodents and bats let their body temperature drop to well below what is typical for most mammals.



Most birds and mammals maintain a body temperature that varies less than 2°C: they are true homeotherms.

Poikilothermic


Increasingly homeothermic

Homeothermic


Body temperature varies with the environmental temperature. Traditionally includes all animals other than birds and mammals, but many reptiles, some large insects and some large fish are not true poikilotherms because they may maintain body temperatures that are different from the surrounding environment.

Body temperature remains almost constant despite environmental fluctuations. Traditionally includes birds and mammals, which typically maintain body temperatures close to 37-38°C. Many reptiles are partially homeothermic and achieve often quite constant body temperatures through behavioral mechanisms.

Source of Body Heat




With a few exceptions, most fish are fully ectothermic. Unlike many reptiles they do not usually thermoregulate.



Snakes use heat energy from the environment to increase their body temperature for activity.



Some large insects like bumblebees may raise their temperature for short periods through muscular activity.



Mammals (and birds) achieve high body temperatures through metabolic activity and reduction of heat losses.

Ectothermic

Increasingly endothermic

Endothermic


Ectotherms depend on the environment for their heat energy. The term ectotherm is often equated with poikilotherm, although they are not the same. Poikilotherms are also ectotherms but many ectotherms may regulate body temperature (often within narrow limits) by changing their behavior (e.g. snakes and lizards).

Endotherms rely largely on metabolic activity for their heat energy. Since they usually maintain a constant body temperature, most endotherms are also homeotherms. As well as birds and mammals, some fast swimming fish, like tuna, and some large insects may also use muscular activity to maintain a high body temperature.

Daily temperature variations in ectotherms and endotherms

Ectotherm: Diurnal lizard (top right)
Body temperature is regulated by behavior so that it does not rise above 40°C. Basking increases heat uptake from the sun. Activity occurs when body temperature is high. Underground burrows are used for retreat.

Endotherm: Human (bottom right)
Body temperature fluctuates within narrow limits over a 24 hour period. Exercise and eating increase body temperature for a short time. Body temperature falls during rest and is partly controlled by an internal rhythm.




Explain why the term "poikilotherm" is not a good term for classifying many terrestrial lizards and snakes:

2. Ectotherms will often maintain high, relatively constant body temperatures for periods in spite of environmental fluctuations, yet they also tolerate marked declines in body temperature to levels lower than are tolerated by endotherms.

(a) Describe the advantages of letting body temperature fluctuate with the environment (particularly at low temperature):

(b) Suggest why ectothermy is regarded as an adaptation to low or variable food supplies:

3. Some endotherms do not always maintain a high body temperature. Some, such as small rodents, allow their body temperatures to fall during hibernation. Explain the advantage of this behavior:



4. The two graphs above illustrate the differences in temperature regulation between a homeothermic endotherm and a poikilothermic ectotherm (such as a fish). Graph A shows change in body temperature with environmental temperature. Graph B shows change in oxygen consumption with environmental temperature. Use the graphs to answer the following:

(a) Explain how ectotherms and endotherms differ in their response to changes in environmental temperature (graph A):

(b) Explain why a poikilothermic ectotherm (no behavioral regulation of temperature) would be limited to environments where temperatures were below about 40°C:

(c) In graph B, state the optimum temperature range for an endotherm:

(d) For an endotherm, the energetic costs of temperature regulation (as measured by oxygen consumption) increase markedly below about 15°C and above 35°C. Explain why this is the case:

(e) For an ectotherm (Graph B), energy costs increase steadily as environmental temperature increases. Explain why: